


Why is VIIRS Ice Surface Temperature Important?

Ice surface temperature (IST) is the radiating, or “skin” temperature at the ice surface. It includes the aggregate temperature of objects comprising the ice surface, such as snow and melt ponds. IST indicates the energy balance at the ice surface, which affects the energy exchange between the ice layer and atmosphere. The VIIRS Ice Surface Temperature algorithm works on clear (non-cloudy) pixels over frozen lakes or rivers as well as the ocean, but not over land ice.



Determination of Ice Surface Temperature:

Ice surface temperature is retrieved using brightness temperatures at split window channels at 11 and 12 μm, which correspond to the M15 and M16 bands for VIIRS, and the satellite sensor scan angle. The algorithm is from the work of Key et al. (1997). The following equation is used:

$$T_s = a + bT_{11} + cT_{12} + d [(T_{11} - T_{12})(\sec\theta - 1)]$$

where T_s is the estimated surface skin temperature, T_{11} and T_{12} are the brightness temperatures at 11 and 12 μm, and θ is the sensor scan angle. The coefficients a , b , c , and d are each derived separately for the Arctic and Antarctic.

Impact on Operations

Primary Application: Satellite retrievals of IST have coverage that far surpasses former methods by ship, buoy, or ice camp observations. Unprecedented coverage allows for the improved understanding of ice surface melt and morphology.

Application: Long-term records of IST can be used to detect and understand the greenhouse effect and climate change in polar regions.

Limitations

Limitation: Algorithm depends on cloud-free retrievals. A low-quality cloud mask may result in incorrect surface temperature.

