

# **OBSERVING SYSTEM SIMULATION EXPERIMENTS TO DEFINE LIDAR WIND REQUIREMENTS**

Robert Atlas

Data Assimilation Office, Code 910.3, NASA/Goddard Space Flight Center  
Greenbelt, Maryland 20771

## **1. Introduction**

Since the advent of meteorological satellites in the 1960's, numerous experiments have been conducted in order to evaluate the impact of these and other data on atmospheric analysis and prediction. Such studies have included both OSE'S and OSSE's. The OSE's were conducted to evaluate the impact of specific observations or classes of observations on analyses and forecasts. Such experiments have been performed for selected types of conventional data and for various satellite data sets as they became available. (See for example the 1989 ECMWF/EUMETSAT workshop proceedings on "The use of satellite data in operational numerical weather prediction" and the references contained therein.) The OSSE's were conducted to evaluate the potential for future observing systems to improve Numerical Weather Prediction (NWP) and to plan for the Global Weather Experiment and more recently for EOS ([1], [2], [3]). In addition, OSSE's have been run to evaluate trade-offs in the design of observing systems and observing networks ([4], [5]), and to test new methodology for data assimilation ([6]).

## **2. Experimental Design**

Although there are many possibilities for how an OSE may be conducted, the most typical procedure is as follows: First a "Control" data assimilation cycle is performed. This is followed by one or more experimental assimilations in which a particular type of data (or specific observations) are either withheld or added to the Control. Forecasts are then generated from both the Control and experimental assimilations every few days (to achieve relative independence of the forecast sample). The analyses and forecasts from each assimilation are then verified and compared in order to determine the impact of each data type being evaluated.

Experiments performed in this manner provide a quantitative assessment of the value of a selected type of data to the specific data assimilation system (DAS) that was used. In addition, the OSE also provides useful information on the effectiveness of the DAS. This information can be used to improve the utilization of this and other data in the DAS, as well as to determine the value of the data.

The methodology currently used for OSSE's is very similar to that described above for OSE's. However, this methodology has evolved considerably since these experiments were first carried out in the 1950's and 60's ([2]). The earliest simulation studies proceeded according to the following sequence of steps. First, an artificial history of the atmosphere was created by numerical integration of a model. Second, simulated "data" were created from the history by the addition of random variations to the history values for temperature, wind, and pressure. Third, the numerical integration that created the history was repeated, but with the meteorological variables in the model replaced by the simulated data at locations and times corresponding to the assumed pattern of observations.

OSSE's of this type were conducted by Charney, Halem and Jastrow, Jastrow and Halem, Williamson and Kasahara, Kasahara, Gordon, and others in preparation for the Global Weather Experiment ([7], [8], [9], [10], [11], [12], [13]). These studies provided an analysis of the Global Atmospheric

Research Program (GARP) data requirements, the “useful” range of predictability, and the need for reference level data. From the results, it was concluded that the assimilation of satellite-derived temperature profiles meeting the GARP data specifications would yield a substantial improvement to the accuracy of numerical weather forecasts. A later study by Cane *et al.* using a modified OSSE procedure indicated similar potential for satellite surface wind data [14].

An examination of the underlying rationale for the early simulation studies (such as [10]), as well as a comparison of the results of the above studies with the results of subsequent real data impact tests (*e.g.* [15] and [16]) indicated several important limitations. The most important weakness stems from the fact that the same numerical model was used both to generate the simulated observations and to test the effectiveness of these observations. (This is referred to as the “identical twin” problem.) Other weaknesses relate to the model-dependence of the studies and the specification of observational errors as random. These weaknesses can lead to overly optimistic results and incorrect conclusions from an OSSE.

The current methodology used for OSSE’s was designed to increase the realism and usefulness of such experiments [17]. In essence, the analysis/forecast simulation system consists of the following elements :

- 1) A long atmospheric model integration using a very high resolution “state of the art” numerical model to provide a complete record of the assumed “true” state of the atmosphere (referred to as the “nature run” or “reference atmosphere”). For the OSSE to be meaningful, it is essential that the nature run be realistic, *i.e.* possess a model climatology, average storm tracks, etc. that agrees with observations to within prespecified limits.
- 2) Simulated conventional and space-based observations from the nature run. All of the observations should be simulated with observed (or expected) coverages, resolution, and accuracy. In addition, bias and horizontal and vertical correlations of errors with each other and with the synoptic situation should be introduced appropriately. Two approaches have been used for this purpose ([17], [3]). The simpler approach is to interpolate the nature run values to the observation locations and then add appropriate errors. The more complicated (and expensive) approach is to attempt to retrieve observations from the nature run in the same way as observations are retrieved from the real atmosphere.
- 3) Control and experimental data assimilation cycles. These are identical to the assimilation cycles in an OSE except that only simulated data are assimilated. In order to avoid the identical twin problem, a different model from that used to generate the nature run is used for assimilation and forecasting. Typically this model has less accuracy and resolution than the nature model. Ideally, the differences between the assimilation and nature models should approximate the differences between a “state of the art” model and the real atmosphere.
- 4) Forecasts produced from the Control and Experimental assimilations. As with the OSE’s, forecasts are generated every few days to develop an independent sample. The analyses and forecasts are then verified against the nature run to obtain a quantitative estimate of the impact of proposed observing systems and the expected accuracies of the analysis and forecast products that incorporate the new data.

An important component of the OSSE that improves the interpretation of results is validation against a corresponding OSE. In this regard, the accuracy of analyses and forecasts and the impact of already existing observing systems in simulation is compared with the corresponding accuracies and data impacts in the real world. Ideally, both the simulated and real results should be similar. Under these conditions, no calibration is necessary and the OSSE results may be interpreted directly. If this is not the case, then calibration of the OSSE results can be attempted by determining the constant of proportionality between the OSE and OSSE impact as described in [3].

### 3. Results and Conclusions

A large number of observing system experiments (OSE's) have been conducted to assess the impact of different types of atmospheric observations. In general, these experiments have shown space-based observing systems to be very important components of the Global Observing System. Nevertheless, conventional surface-based observations are still of critical importance.

Observing system simulation experiments (OSSE's) provide an effective means to evaluate the potential impact of a proposed observing system, as well as to determine tradeoffs in their design, and to evaluate data assimilation methodology. Great care must be taken to ensure realism of the OSSE's, and in the interpretation of OSSE results. OSSE's have shown considerable potential for space-based lidar wind profiles and for improved space-based surface winds. However, quantitative results from OSSE's may only be valid for instruments meeting the expected accuracies and coverages that were simulated, and for the data assimilation systems used in the particular OSSE.

At the conference, results will be presented from OSSE's that have been conducted over the last several years at the DAO. These will include OSSE's aimed at determining the requirements for and assessing the potential impact of space-based lidar winds and other advanced sounders, as well as experiments to evaluate trade-offs in the design of future observing systems.

#### ACKNOWLEDGMENTS

The author gratefully acknowledges the support of NASA HQ . Several of the OSSE's referred to in this paper were performed in collaboration with G.D. Emmitt and S. Wood. I would also like to thank S.C. Bloom, J.C. Jusem, E. Brin, J. Ardizzone, J.Terry, and D. Bungato for significant contributions to this work.

#### REFERENCES

- [1] E. Kalnay R. Atlas and M. Halem, "The impact of satellite temperature sounding and wind data on numerical weather prediction," *Optical Engineering*, vol. 24, pp. 341–346, 1985.
- [2] C.P. Arnold jr. and C.H. Dey, "Observing-system simulation experiments: Past, present, and future," *Bull. Amer. Meteor. Soc.*, vol. 67, pp. 687–695, 1986.
- [3] R. Isaacs J. Louis T. Nehr Korn R.N. Hoffman, C.Grassotti and D. Norquist, "Assessment of the impact of simulated satellite lidar wind and retrieved 183 ghz water vapor observations on a global data assimilation system," *Mon. Wea. Rev.*, vol. 118, pp. 490–513, 1990.
- [4] R. Atlas and G.D. Emmitt, "Implications of several orbit in-clinations for the impact of laws on global climate studies," in *Second Symposium on Global Change Studies*, New Orleans, 1991, pp. 28–32.
- [5] G.D. Rohaly and T.N. Krishnamurti, "An observing system simulation experiment for the laser atmospheric wind sounder (laws)," *J. Applied Meteor.*, vol. 32, pp. 1453–1471, 1993.
- [6] R. Atlas and S. Bloom, "Global surface wind vectors resulting from the assimilation of satellite wind speed data in atmospheric general circulation models," in *Oceans 89 Proceedings*. 1989, pp. 260–265, IEEE.
- [7] M. Halem J. Charney and R. Jastrow, "Use of incomplete historical data to infer the present state of the atmosphere," *J. Atmos. Sci.*, vol. 26, pp. 1160–1163, 1969.

- [8] M. Halem and R. Jastrow, "Analysis of garp data requirements," *J. Atmos. Sci.*, vol. 27, pp. 177, 1970.
- [9] R. Jastrow and M. Halem, "Simulation studies related to garp," *Bull. Amer. Meteor. Soc.*, vol. 51, pp. 1705–1708, 1970.
- [10] R. Jastrow and M. Halem, "Accuracy and coverage of temperature data derived from the ir radiometer on the NOAA 2 satellite," *J. Atmos. Sci.*, vol. 30, pp. 958–964, 1973.
- [11] D. Williamson and A. Kasahara, "Adaptation of meteorological variables forced by updating," *J. Atmos. Sci.*, vol. 28, pp. 1313–1324, 1971.
- [12] Kasahara, "Simulation experiments for meteorological observing systems for garp," *Bull. Amer. Meteor. Soc.*, vol. 53, 1972.
- [13] Jr. C.T. Gordon, L. Umscheid and K. Miyakoda, "Simulation experiments for determining wind data requirements in the tropics," *J. Atmos. Sci.*, vol. 29, pp. 1064–1075, 1972.
- [14] M. Halem M.A. Cane, V.J. Cardone and I. Halberstam, "On the sensitivity of numerical weather prediction to remotely sensed marine surface wind data: A simulation study," *J. Geophys. Res.*, vol. 86, pp. 8093–8106, 1981.
- [15] W. Baker M. Halem, E. Kalnay-Rivas and R. Atlas, "An assessment of the fgge satellite observing systems during sop-1," *Bull. Amer. Meteor. Soc.*, vol. 63, pp. 407–426, 1982.
- [16] W.E. Baker, G.D. Emmitt, F. Robertson, R.M. Atlas, J.E. Molinari, D.A. Bowdle, J. Paegle, R.M. Hardesty, R.T. Menzies, T.N. Krishnamurti, R.A. Brown, M.J. Post, J.R. Anderson A.C. Lorenc, and J. McElroy, "Lidar measured winds from space: A key component for weather and climate prediction," *Bull. Amer. Meteor. Soc.*, vol. 76, no. 6, pp. 869–888, 1995.
- [17] J. Susskind W. E. Baker R. Atlas, E. Kalnay and M. Halem, "Simulation studies of the impact of future observing systems on weather prediction," in *Proc. Seventh Conf. on NWP*, 1985, pp. 145–151.