

# Assimilation of AMSU-A in the presence of cloud and precipitation



Peter Weston, Alan Geer and Niels Bormann

Research Department, ECMWF, Reading, United Kingdom

## 1) Introduction

In recent years a number of NWP centres have found significant positive impact from assimilating microwave radiances in all-sky conditions (Geer et al., 2017, Zhu et al., 2016, Migliorini and Candy, 2019). The clear-sky assimilation of AMSU-A at ECMWF has been fine-tuned over the past two decades, but the cloud detection still rejects up to 25% of data from the tropospheric temperature sounding channels. Assimilating the additional cloud- and precipitation-affected data should allow some positive forecast

impact. At ECMWF two largely different systems and code paths are used for clear-sky and all-sky assimilation. Many of the differences between the two systems have now been eliminated but matching the performance of the existing clear-sky assimilation of AMSU-A in the all-sky system has proved challenging. This poster highlights remaining differences between the clear- and all-sky systems, compares performance of AMSU-A clear- and all-sky assimilation and suggests potential future enhancements. For more details see Weston et al. (2019).

## 2) All-sky v clear-sky differences

Aspect	Clear-sky	All-sky
Radiative transfer	RTTOV	RTTOV-SCATT
Cloud detection	On	Off
Observation errors	Noise, cloud and surface dependent	Noise and cloud dependent
Thinning	125km x 125km boxes	Alternate points of a $T_L255$ reduced Gaussian grid, threshold on distance from grid point
Skin temperature sink variable	On	Off
Quality control	Basic, fg_depar, VarQC	Additional snow/sea-ice screening

Table 1: Differences between clear- and all-sky assimilation configurations. Aspects not covered such as bias correction, surface emissivity treatment etc. are the same in clear- and all-sky systems.

The following assimilation choices were important to achieve the current performance: thinning all AMSU-A's from different satellites together rather than separately; additional snow/sea-ice screening due to residual biases; matching up the bias predictors and initial coefficients; changing the interpolation method from nearest neighbour to bi-linear in all-sky. These changes led to improved first guess fits to observations, especially ATMS, and improved forecast scores, particularly at high latitudes.

## 4) Possible future enhancements




Figure 4: Standard deviation of first guess departures for AMSU-A channel 5 binned by cloud liquid water path for the tropics (upper) and mid-latitudes (lower).

Larger cloudy first guess departures in tropics (e.g. from convection mislocation). This could be accounted for using an additional total column water vapour predictor in the observation error model.

More observations assimilated at edge of scan in clear-sky. All-sky thinning could be done on a  $T_L159$  grid and distance threshold relaxed to better match the clear-sky observation distribution





Figure 5: Number of assimilated observation counts across the scanline for AMSU-A all-sky and clear-sky.

## 3) Results v denial



AMSU-A all-sky and clear-sky result in similar (large) improvements to extra-tropical geopotential height forecasts against an AMSU-A denial.

Figure 1 (above): Change in RMSE of extra-tropical 500hPa geopotential height forecasts for AMSU-A all-sky (black) and clear-sky (red) against an AMSU-A denial baseline

All-sky improves fits to wind observations slightly more than clear-sky in the extra-tropics.

All-sky improves fits to humidity sensitive observations (AMSR2 and ATMS channels 18-22) more than clear-sky in the extra-tropics.

All-sky and clear-sky have similar impacts to fits to temperature sensitive observations (ATMS channels 6-15 and GPSRO).





Figure 2 (above): Change in first guess fits to AMSR2 (upper), conventional wind (lower left), and GPSRO (lower right) observations in the Southern hemisphere for AMSU-A all-sky (black) and clear-sky (red) against an AMSU-A denial baseline.

Figure 3 (below): Change in RMSE of tropical 850hPa vector wind (left) and temperature (right) forecasts for AMSU-A all-sky (black) and clear-sky (red) against an AMSU-A denial baseline.



All-sky degrades short-range wind and temperature forecasts in the tropics compared to clear-sky.

## 5) Conclusions

In the extra-tropics, the performance of AMSU-A all-sky assimilation is now comparable to the existing clear-sky assimilation. This has involved bringing many aspects of the all-sky assimilation configuration into line with the clear-sky system e.g. thinning, bias correction, quality control. Some of these aspects had surprisingly large impacts on forecast accuracy.

However, in the tropics, AMSU-A all-sky degrades forecasts when compared to the existing clear-sky assimilation. We have a couple of hypotheses for why this might be the case: larger first guess departures than assigned observation errors in cloudy areas in the tropics and residual biases over arid regions. In addition, the different distribution of observations across scan lines could also be sub-optimal globally.

Future work will aim to address these issues by accounting for additional observation error variation using an additional predictor such as total column water vapour, eliminating the remaining thinning differences and additional quality control over arid areas. With these enhancements we aim to make the all-sky AMSU-A assimilation ready for operational implementation.

## References

Geer, A. J., Baordo, F., Bormann, N., Chambon, P., English, S. J., Kazumori, M., Lawrence, H., Lean, P., Lonitz, K., Lupu, C., 2017. The growing impact of satellite observations sensitive to humidity, cloud and precipitation. Q.J.R. Meteorol. Soc. 143 (709), 3189–3206. doi: 10.1002/qj.3172

Migliorini, S., Candy, B., 2019. All-sky satellite data assimilation of microwave temperature sounding channels at the Met Office. Quarterly Journal of the Royal Meteorological Society 145 (719), 867–883. doi: 10.1002/qj.3470

Weston, P., Geer, A., Bormann, N. 2019. Investigations into the assimilation of AMSU-A in the presence of cloud and precipitation. EUMETSAT/ECMWF Fellowship Programme Research Report No.50

Zhu, Y., Liu, E., Mahajan, R., Thomas, C., Groff, D., Van Delst, P., Collard, A., Kleist, D., Treadon, R., Derber, J. C., 2016. All-sky microwave radiance assimilation in NCEP's GSI analysis system. Monthly Weather Review 144 (12), 4709–4735. doi: 10.1175/MWR-D-15-0445.1

## Acknowledgments

Peter Weston was funded by the EUMETSAT Fellowship programme